arXiv:1502.03266v3 [math.PR] 14 Sep 2016

Laplace approximation with estimated error and application to probability

Vassili Kolokoltsov (v.kolokoltsov@warwick.ac.uk),
Tomasz M. Lapinski (84tomek@gmail.com)

University of Warwick,
Department of Statistics,
CV4 7AL Coventry, UK,

September 15, 2016

Abstract

In this paper we prove a Multivariate Laplace approximation with estimated error for two cases of maximum, in the interior of the domain and on the boundary. Additionally, the function in the exponent and its point of maximum depends on the integrals asymptotic parameter. As an application, we prove Law of Large Numbers and the Central Limit Theorem for the random vector related to the Laplace Integral. The second result gives different limiting distribution for two cases of maximum in the Laplace integral. When it is in the interior of the domain it is Gaussian distribution and on the boundary it is exponential in one direction and Gaussian in other directions.

Key words— Multivariate Laplace method, Laplace method, Asymptotic approximation of integrals, Law of Large Numbers, Central Limit Theorem

1 Introduction

We prove Multivariate Laplace approximation which is an extension of results already known in the literature, see for example Wong [2001],Breitung [1994]. Additionally, we consider the function in the exponent of the integral and its point of maximum dependent on 'large' parameter. We also provide rate of convergence and estimate of the remainder term of developed approximations. These results are development of results in Kolokoltsov [2000]. Moreover, we consider two cases of maximum, in the interior and on the boundary of the domain.

For the probability application we first define a probability space, with domain of Laplace integral as a set and Borel σ-algebra on it. The distribution of particular event is constructed by taking the Laplace integral over the event set and normalizing by integral over the domain. We consider a random vector whose values are the points of the domain of considered integral. The first probabilistic result, Law of Large Numbers gives us mean equal to the point of maximum of the function in the exponent of Laplace integral. The second result, Central Limit Theorem gives us the distribution of fluctuations different for two cases of maximum. When it is in the interior of the domain it is Gaussian and on the boundary, exponential in one direction and Gaussian in other directions. Additionally, rate of convergence for both limit theorems is included. In the proof we used developed Laplace approximation and some rather standard estimates.

The paper consists of three sections and appendix. Second section contains our main results, Multidimensional Laplace approximation. First we consider one dimensional case with maximum on the boundary. Then we consider multidimensional case with the maximum in the interior and then general case on the boundary. In the third section we develop the probability application. We introduce the probability space and random vectors, state and prove two limit theorems. In the end of this section we include additional estimates used in the proofs of the probability results. In the appendix we included some standard, easily obtained results used throughout the paper.

2 Laplace approximation

We consider an integral over a bounded domain $\Omega \subset \mathbb{R}^m$

$$I(N) := \int_\Omega g(x) e^{N f(x,N)} dx, \tag{1}$$

where $N > N_0$, for some $N_0 \in \mathbb{N}$, are positive integers.
Function $f : \Omega \times \mathbb{N} \to \mathbb{R}$ converges to $\mathbb{R}$-valued function as $N \to \infty$ and have a unique maximum in the points denoted $x^*(N)$ for all $N > N_0$ and x^* in the limit. Additionally, it is three times continuously differentiable in some closed neighborhood Ω' of x^* and those derivatives are uniformly convergent as $N \to \infty$.
Further, we consider two types of maximum of f:

(a) in the interior of Ω and x^* is a non-degenerate critical point,

(b) on the boundary $\partial\Omega$ where x^* is a non-degenerate critical point and non-critical in the orthogonal direction. Here we assume $\partial\Omega$ is a hyperplane in some neighborhood of x^*.

Function $g : \Omega \to \mathbb{R}$ is continuously differentiable in Ω' and bounded in Ω.

From assumptions about f and g and for appropriately chosen Ω' we have

$$F^{(2)} := \sup_{x \in \Omega', N > N_0} \|D^2 f(x,N)\| < \infty, \tag{2}$$

$$F'^{(2)} := \inf_{x \in \Omega', N > N_0} \|D^2 f(x,N)^{-1}\|^{-1} > 0, \tag{3}$$

$$F'^{(2)}_\Omega := \min\left\{ F'^{(2)}, \inf_{N > N_0} \frac{f(x^*(N), N) - \sup_{x \in \Omega \setminus \Omega'} f(x,N)}{\sup_{x \in \Omega \setminus \Omega'} |x - x^*(N)|^2} \right\} > 0, \tag{4}$$

$$F^{(3)} := \sup_{x \in \Omega', N > N_0} \|D^{\otimes 3} f(x,N)\| < \infty, \tag{5}$$

$$G := \sup_{x \in \Omega} \|g(x)\| < \infty, \tag{6}$$

$$G^{(1)} := \sup_{x \in \Omega'} \|Dg(x)\| < \infty. \tag{7}$$

Additionally, we choose Ω' and N_0 such that for all $N > N_0$ the points $x^*(N) \in \Omega'$ and for all $x \in \Omega'$,

$$\lambda \leq |\det(D^2 f(x,N))| \leq \Lambda, \tag{8}$$

where λ and Λ are some positive constants.
For the maximum of type (b) the derivatives in above determinant and those in the constant $F'^{(2)}$ are w.r.t. coordinates of the hyperplane $\partial\Omega$ at x^* and also only for this maximum we define

$$F'^{(1)} := \inf_{x\in\Omega', N>N_0} |f'(x,N)| > 0, \tag{9}$$

$$F'^{(1)}_{\Omega} := \min\left\{F'^{(1)}, \inf_{N>N_0} \frac{f(x^*(N),N) - \sup_{x\in\Omega\setminus\Omega'} f(x,N)}{\sup_{x\in\Omega\setminus\Omega'} |x - x^*(N)|}\right\} > 0, \tag{10}$$

where the derivative is in direction orthogonal to that hyperplane.

2.1 One-dimensional domain with the maximum on the boundary

Theorem 1. *For the integral (1) with $\Omega \subset \mathbb{R}$ and the maximum of type (b), we have following approximation*

$$\int_\Omega g(x) e^{Nf(x,N)} dx = e^{Nf(x^*,N)} \frac{1}{N}\left(\frac{g(x^*)}{|f'(x^*,N)|} + \frac{\omega_B(N)}{N}\right),$$

where $\omega_B(N)$ is some bounded function and

$$|\omega_B(N)| \leq \frac{|g(x^*)|F^{(2)}}{\left(F'^{(1)}\right)^3} e^{\frac{1}{2}F^{(2)}} + \frac{G^{(1)}}{\left(F'^{(1)}\right)^2} + \frac{G}{F'^{(1)}_\Omega} N e^{-N^{1/2}F'^{(1)}_\Omega} + \frac{|g(x^*)|}{F'^{(1)}} N e^{-N^{1/2}F'^{(1)}}.$$

Proof. Let us define

$$I_G(N) := g(x^*) e^{Nf(x^*,N)} \frac{1}{N} \frac{1}{|f'(x^*,N)|} = g(x^*) e^{Nf(x^*,N)} \int_{x^*}^{\infty} e^{-N|f'(x^*,N)|(x-x^*)} dx,$$

and introduce the set $U_N := \{x : |x - x^*| \leq 1/N^{1/2},\ N > N_0\}$, with N_0 large enough that $U_N \subset \Omega'$. Then by separating the set U_N and with use of Taylor's Theorem we decompose $I(N)$ and $I_G(N)$

$$\begin{aligned} I(N) =& I_{11}(N) + I_{12}(N) + I_2(N) := g(x^*) \int_{U_N} e^{Nf(x,N)} dx + \int_{U_N} g'(x_{\theta_1})(x - x^*) e^{Nf(x,N)} dx + \\ &+ \int_{\Omega\setminus U_N} g(x) e^{Nf(x,N)} dx, \\ I_G(N) =& I_{G1}(N) + I_{G2}(N) := g(x^*) e^{Nf(x^*,N)} \int_{U_N} e^{-N|f'(x^*,N)|(x-x^*)} dx + \\ &+ g(x^*) e^{Nf(x^*,N)} \int_{\Omega\setminus U_N} e^{-N|f'(x^*,N)|(x-x^*)} dx. \end{aligned}$$

Now, we put together above decompositions

$$|I(N) - I_G(N)| \leq |I_{11}(N) - I_{G1}(N)| + |I_{12}(N)| + |I_2(N)| + |I_{G2}(N)|,$$

and approximate each term.
To approximate the expression $|I_{11}(N) - I_{G1}(N)|$ we apply Taylor's theorem for f and inequality $|e^t - 1| \leq |t|e^{|t|}$. Further, without loss of generality we can assume $f'(x^*, N) < 0$ and get

$$|I_{11}(N) - I_{G1}(N)| = \left| g(x^*) \int_{U_N} e^{Nf(x^*,N) - N|f'(x^*,N)|(x-x^*)} \left(e^{\frac{N}{2} f''(x_{\theta_2},N)(x-x^*)^2} - 1 \right) dx \right| \leq$$

$$\leq \left| g(x^*) \int_{U_N} e^{Nf(x^*,N) - N|f'(x^*,N)|(x-x^*)} \frac{N}{2} f''(x_{\theta_2}, N)(x - x^*)^2 e^{\frac{N}{2} f''(x_{\theta_2},N)(x-x^*)^2} dx \right|.$$

Since the integration is over U_N, we have that $|x - x^*| \leq \frac{1}{N^{1/2}}$, hence above expression is bounded by

$$\frac{1}{2} N f''(x_{\theta_2}, N) e^{Nf(x^*,N) + \frac{1}{2} N f''(x_{\theta_2},N)} |g(x^*)| \int_{U_N} |x - x^*|^2 e^{-N|f'(x^*,N)|(x-x^*)} dx,$$

and the integral

$$\int_{U_N} |x - x^*|^2 e^{-N|f'(x^*,N)|(x-x^*)} dx \leq \int_{x^*}^{\infty} |x - x^*|^2 e^{-N|f'(x^*,N)|(x-x^*)} dx = \frac{2}{(N|f'(x^*, N)|)^3}.$$

Hence we have

$$|I_{11}(N) - I_{G1}(N)| \leq \frac{1}{N^2} \frac{F^{(2)}}{|f'(x^*, N)|^3} |g(x^*)| e^{Nf(x^*,N) + \frac{1}{2} F^{(2)}}.$$

For the integral $|I_{12}(N)|$ we use Taylor's Theorem $f(x, N) = f(x^*, N) + f'(x_{\theta_3}(N), N)(x - x^*)$ to get an estimate for f. Since $f(x^*, N)$ is its uniqe maximum and $U_N \subset \Omega'$ we get

$$f(x, N) \leq f(x^*, N) - F'^{(1)} |x - x^*|,$$

where $F'^{(1)}$ is given by (9).
We insert it into I_{12} and as g has bounded derivative in U_N we integrate

$$|I_{12}(N)| \leq e^{Nf(x^*,N)} \int_{x^*}^{\infty} g'(x_{\theta_1})(x - x^*) e^{-NF'^{(1)}|x-x^*|} dx \leq G^{(1)} e^{Nf(x^*,N)} \frac{1}{\left(NF'^{(1)}\right)^2}.$$

For $|I_2(N)|$ we substitute the estimate of f as for the integral I_{12}, but here we have possibly smaller constant $F'^{(1)}_{\Omega}$ defined by (10). Then find an upper bound and integrate

$$|I_2(N)| \leq \int_{x^*}^{\infty} |g(x)| e^{Nf(x^*,N) - NF'^{(1)}_{\Omega}|x-x^*|} dx = G e^{Nf(x^*,N)} \frac{1}{NF'^{(1)}_{\Omega}} e^{-N^{1/2} F'^{(1)}_{\Omega}}.$$

The integral $|I_{G2}(N)|$ we approximate similarly, but instead of $x_{\theta_3}(N)$ we have x^* in the first derivative of Taylor expansion, hence

$$|I_{G2}(N)| \leq |g(x^*)| e^{Nf(x^*,N)}) \frac{1}{N|f'(x^*, N)|} e^{-N^{1/2}|f'(x^*,N)|}.$$

Now, we combine all approximations and conclude with

$$|I(N) - I_G(N)| \leq e^{Nf(x^*,N)} \frac{1}{N^2} \left[\frac{|g(x^*)| F^{(2)}}{\left(F'^{(1)}\right)^3} e^{\frac{1}{2} F^{(2)}} + \frac{G^{(1)}}{\left(F'^{(1)}\right)^2} + \frac{G}{F'^{(1)}_{\Omega}} N e^{-N^{1/2} F'^{(1)}_{\Omega}} + \right.$$

$$\left. + \frac{|g(x^*)|}{F'^{(1)}} N e^{-N^{1/2} F'^{(1)}} \right],$$

and therefore we proved the theorem. ☐

2.2 m-dimensional function with the maximum in the interior of the domain

Theorem 2. *For the integral (1) with $\Omega \subset \mathbb{R}^m$ and the maximum of type (a) we have following approximation*

$$\int_\Omega g(x)e^{Nf(x,N)}dx = e^{Nf(x^*(N),N)}\left(\frac{2\pi}{N}\right)^{\frac{m}{2}}\left(\frac{g(x^*(N))}{\sqrt{\det D^2 f(x^*(N),N)}} + \frac{\omega_I(N)}{\sqrt{N}}\right),$$

where $\omega_I(N)$ is some bounded function and

$$\begin{aligned}|\omega_I(N)| \leq &\pi^{\frac{m}{2}}\frac{\Gamma(\frac{m+1}{2})}{\Gamma(\frac{m}{2})}\left(\frac{F'^{(2)}}{2}\right)^{-\frac{m+1}{2}}\left[\frac{F^{(3)}}{3F'^{(2)}}e^{F^{(3)}}|g(x^*(N))| + G^{(1)}\right] + \frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})}\left(\frac{F_\Omega'^{(2)}}{2}\right)^{-\frac{m+1}{2}} \times \\ &\times \left[\Gamma(m) + \left(1 + \sqrt{F_\Omega'^{(2)}}\frac{N^{1/6}}{\sqrt{2}}\right)^{m-1}\right]\big(G + |g(x^*(N))|\big)e^{-N^{1/3}F_\Omega'^{(2)}}.\end{aligned}$$

Proof. Since f has property (8) we can define

$$\begin{aligned}I_G(N) :&= g(x^*(N))e^{Nf(x^*(N),N)}\left(\frac{2\pi}{N}\right)^{\frac{m}{2}}\frac{1}{\sqrt{\det D^2 f(x^*(N),N)}} = \\ &= g(x^*(N))e^{Nf(x^*(N),N)}\int_{\mathbb{R}^m}\exp\Big(\frac{1}{2}N(x - x^*(N))^T D^2 f(x^*(N),N)(x - x^*(N))\Big)dx,\end{aligned}$$

and introduce the set $U_N = \{x : |x - x^*(N)| \leq \frac{1}{N^{1/3}},\ N > N_0\}$ with N_0 large enough to $U_N \subset \Omega'$. Then by separating the set U_N and using Taylors theorem we decompose $I(N)$ and $I_G(N)$

$$\begin{aligned}I(N) =&I_{11}(N) + I_{12}(N) + I_2(N) := g(x^*(N))\int_{U_N}e^{Nf(x,N)}dx + \\ &+ \int_{U_N}Dg(x_{\theta_1}(N))^T(x - x^*(N))e^{Nf(x,N)}dx + \int_{\Omega\setminus U_N}g(x)e^{Nf(x,N)}dx, \\ I_G(N) =& I_{G1}(N) + I_{G2}(N) := \\ &= g(x^*(N))e^{Nf(x^*(N),N)}\int_{U_N}\exp\Big(\frac{1}{2}N(x - x^*(N))^T D^2 f(x^*(N),N)(x - x^*(N))\Big)dx + \\ &+ g(x^*(N))e^{Nf(x^*(N),N)}\int_{\mathbb{R}^m\setminus U_N}\exp\Big(\frac{1}{2}N(x - x^*(N))^T D^2 f(x^*(N),N)(x - x^*(N))\Big)dx.\end{aligned}$$

We combine above integrals into

$$|I(N) - I_G(N)| \leq |I_{11}(N) - I_{G1}(N)| + |I_{12}(N)| + |I_2(N)| + |I_{G2}(N)|.$$

For the expression $I_{11}(N) - I_{G1}(N)$ we use third order Taylor's theorem to obtain

$$\begin{aligned}&|I_{11}(N) - I_{G1}(N)| = \\ &= |g(x^*(N))|\int_{U_N}\exp\Big(Nf(x^*(N),N) + \frac{N}{2}(x - x^*(N))^T D^2 f(x^*(N),N)(x - x^*(N))\Big)\times \\ &\times \Big(\exp\Big(\frac{1}{6}ND^{\otimes 3}f(x_{\theta_2}(N),N)(x - x^*(N))^{\otimes 3}\Big) - 1\Big)dx,\end{aligned}$$

where $Df(x^*(N),N)^T(x-x^*(N))=0$ for all x since $x^*(N)$ is a critical point.
Next, we apply inequality $|e^t-1|\le |t|e^{|t|}$ and use an upper bound $\left|D^{\otimes 3}f(x_{\theta_2}(N),N)(x-x^*(N))^{\otimes 3}\right|\le F^{(3)}|x-x^*(N)|^3$ derived in (17) in Appendix and since $|x-x^*(N)|\le \frac{1}{N^{1/3}}$ for $x\in U_N$ we conclude

$$\begin{aligned}&|I_{11}(N)-I_{G1}(N)|\le\\ &\quad\le \frac{1}{6}NF^{(3)}e^{Nf(x^*(N),N)+F^{(3)}}|g(x^*(N))|\int_{U_N}|x-x^*(N)|^3e^{\frac{N}{2}(x-x^*(N))^TD^2f(x^*(N),N)(x-x^*(N))}dx.\end{aligned}$$

Then using calculus result $\int_{\mathbb{R}^m}|x|^3e^{-N|x|^2}dy=N^{-\frac{m+3}{2}}\pi^{n/2}\Gamma(\frac{m+1}{2})/\Gamma(\frac{m}{2})$ we obtain the estimate

$$\begin{aligned}&\int_{U_N}|x-x^*(N)|^3e^{\frac{N}{2}(x-x^*(N))^TD^2f(x^*(N),N)(x-x^*(N))}dx\le\int_{\mathbb{R}^m}|x|^3e^{-\frac{1}{2}NF'^{(2)}x^2}dx=\\ &=\pi^{\frac{m}{2}}\frac{\Gamma(\frac{m+1}{2})}{\Gamma(\frac{m}{2})}\left(\frac{1}{2}NF'^{(2)}\right)^{-\frac{m+3}{2}},\end{aligned}$$

and therefore

$$|I_{11}(N)-I_{G1}(N)|\le\frac{1}{6}F^{(3)}e^{F^{(3)}}\pi^{m/2}\frac{\Gamma(\frac{m+1}{2})}{\Gamma(\frac{m}{2})}\left(\frac{F'^{(2)}}{2}\right)^{-\frac{m+3}{2}}|g(x^*(N))|\frac{1}{\sqrt{N}}e^{Nf(x^*(N),N)}N^{-\frac{m}{2}}.$$

For the integral $I_{12}(N)$, by Taylor's Theorem and taking into account that $|x-x^*|\|D^2f(x_\theta)^{-1}\|^{-1}\le\left|D^2f(x_\theta)^T(x-x^*)\right|$ we have an upper bound of $f(x,N)$ for any $x\in U_N$

$$f(x,N)\le f(x^*(N),N)-\frac{1}{2}F'^{(2)}(x-x^*(N))^2,$$

and since g has bounded derivative in U_N we have

$$|I_{12}(N)|\le G^{(1)}e^{Nf(x^*(N),N)}\int_{U_N}|x-x^*(N)|e^{-\frac{1}{2}NF'^{(2)}(x-x^*(N))^2}dx,$$

and with use of $\int_{\mathbb{R}^m}|x|e^{-N|x|^2}dy=N^{-\frac{m}{2}}\pi^{m/2}\Gamma(\frac{m+1}{2})/\Gamma(\frac{m}{2})$ we obtain

$$|I_{12}(N)|\le\pi^{\frac{m}{2}}\frac{\Gamma(\frac{m+1}{2})}{\Gamma(\frac{m}{2})}\left(\frac{F'^{(2)}}{2}\right)^{-\frac{m+1}{2}}G^{(1)}\frac{1}{\sqrt{N}}e^{Nf(x^*(N),N)}N^{-\frac{m}{2}}.$$

For $I_2(N)$, since Ω is bounded we have the same estimate for f as in I_{12} but with constant $F_\Omega'^{(2)}$

$$|I_2(N)|\le e^{Nf(x^*(N),N)}\int_{\mathbb{R}^m\setminus U_N}g(x)e^{-\frac{1}{2}NF_\Omega'^{(2)}(x-x^*(N))^2}dx,$$

and since g is bounded we estimate above using (18) from Appendix

$$\begin{aligned}|I_2(N)|\le&\frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})}\left(\frac{NF_\Omega'^{(2)}}{2}\right)^{-\frac{m+1}{2}}\left[\Gamma(m)+\left(1+\sqrt{F_\Omega'^{(2)}}\frac{N^{1/6}}{\sqrt{2}}\right)^{m-1}\right]\times\\ &\times Ge^{Nf(x^*(N),N)}e^{-N^{1/3}F_\Omega'^{(2)}}.\end{aligned}$$

In the case of I_{G2} the approximation procedure is analogical to I_2, yielding

$$\begin{aligned}|I_{G2}(N)| \leq & \frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})}\left(\frac{NF_\Omega'^{(2)}}{2}\right)^{-\frac{m+1}{2}}\left[\Gamma(m)+\left(1+\sqrt{F_\Omega'^{(2)}}\frac{N^{1/6}}{\sqrt{2}}\right)^{m-1}\right]\times \\ & \times g(x^*(N))e^{Nf(x^*(N),N)}e^{-N^{1/3}F_\Omega'^{(2)}}.\end{aligned}$$

Then we combine above approximations into

$$\begin{aligned}|I(N)-I_G(N)| \leq & \frac{1}{\sqrt{N}}e^{Nf(x^*(N),N)}\left(\frac{2\pi}{N}\right)^{\frac{m}{2}}\left[\pi^{\frac{m}{2}}\frac{\Gamma(\frac{m+1}{2})}{\Gamma(\frac{m}{2})}\left(\frac{F'^{(2)}}{2}\right)^{-\frac{m+1}{2}}\times\right. \\ & \times\left(\frac{F^{(3)}}{3F'^{(2)}}e^{F^{(3)}}|g(x^*(N))|+G^{(1)}\right)+\frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})}\left(\frac{F_\Omega'^{(2)}}{2}\right)^{-\frac{m+1}{2}}\left[\Gamma(m)+\right. \\ & \left.\left.+\left(1+\sqrt{F_\Omega'^{(2)}}\frac{N^{1/6}}{\sqrt{2}}\right)^{m-1}\right]\big(G+|g(x^*(N))|\big)e^{-N^{1/3}F_\Omega'^{(2)}}\right],\end{aligned}$$

which yields the statement of the theorem. □

2.3 m-dimensional function with the maximum on the boundary of the domain

Theorem 3. *We consider an integral (1) with $\Omega \subset \mathbb{R}^m$ and with maximum of type (b). If the $\partial\Omega$ at the maximum is not orthogonal to first coordinate axis then we obtain it by appropriate rotation of coordinate system. Then we have following approximation*

$$I(N) = e^{Nf(x^*(N),N)}\frac{1}{N}\left(\frac{2\pi}{N}\right)^{\frac{m-1}{2}}\left(\frac{g(x^*(N))}{|f'(x^*(N),N)|\sqrt{|\det D^2f(x^*(N),N)|}}+\frac{\omega(N)}{\sqrt{N}}\right),$$

where f' is the derivative w.r.t. first coordinate and D^2f is a matrix of second derivatives w.r.t. all others coordinates, $\omega(N)$ is some bounded function and

$$\begin{aligned}|\omega(N)| \leq & \frac{|\omega_{B1}(N)|}{\sqrt{N}}+|\omega_I(N)|\left(\frac{1}{F'^{(1)}}+\frac{|\omega_{B2}(N)|}{N}\right)+\frac{2\sqrt{\pi}}{\Gamma(\frac{m}{2})}\big(F_\Omega'^{(2)}\big)^{-\frac{m+1}{2}}\times \\ & \times\left[\Gamma(m)+\left(1+\sqrt{NF_\Omega'^{(2)}}\frac{\min_{x\in\Omega\setminus\Omega'}|x-x^*|}{\sqrt{2}}\right)^{m-1}\right]Ge^{-NF_\Omega'^{(2)}(\min_{x\in\Omega\setminus\Omega'}|x-x^*|)^2}, \\ |\omega_{B1}(N)| \leq & \frac{|g(x^*(N))|F^{(2)}}{\sqrt{\lambda}\big(F'^{(1)}\big)^3}e^{\frac{1}{2}F^{(2)}}+\frac{1}{\big(F'^{(1)}\big)^2}\left(\frac{G^{(1)}}{\sqrt{\lambda}}+\frac{mGF^{(3)}}{\lambda^{3/2}F'^{(2)}}\right)+\frac{G}{\sqrt{\lambda}F_\Omega'^{(1)}}Ne^{-N^{1/2}F_\Omega'^{(1)}}+ \\ & +\frac{|g(x^*(N))|}{\sqrt{\lambda}F'^{(1)}}Ne^{-N^{1/2}F'^{(1)}}, \\ |\omega_{B2}(N)| \leq & \frac{F^{(2)}}{\big(F'^{(1)}\big)^3}e^{\frac{1}{2}F^{(2)}}+\frac{1}{F_\Omega'^{(1)}}Ne^{-N^{1/2}F_\Omega'^{(1)}}+\frac{1}{F'^{(1)}}Ne^{-N^{1/2}F'^{(1)}}.\end{aligned}$$

where ω_I is a reminder as in the Theorem 2.

Proof. We decompose $I(N)$ into

$$I(N) = I'(N)+I''(N) := \int_{\Omega'}g(x)e^{Nf(x,N)}dx+\int_{\Omega/\Omega'}g(x)e^{Nf(x,N)}dx.$$

Then we approximate $I''(N)$ as the integral I_2 in the proof of the previous theorem but with an integral over domain $\Omega\backslash\Omega'$ instead of $\Omega\backslash U_N$. Therefore we use (19) and obtain

$$|I''(N)| \leq \frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})}\left(\frac{NF_{\Omega}'^{(2)}}{2}\right)^{-\frac{m+1}{2}}\left[\Gamma(m)+\left(1+\sqrt{NF_{\Omega}'^{(2)}}\frac{R}{\sqrt{2}}\right)^{m-1}\right]Ge^{Nf(x^*(N),N)}e^{-NF_{\Omega}'^{(2)}R^2},$$

where $R = \min_{x\in\Omega\backslash\Omega'}|x-x^*|$.
Let us define $\hat{x} := (x_2, x_3, \ldots, x_m), x = (x_1, \hat{x})$ and the curve of points of maximal values of f along variable x_1 as $\hat{x}^*(N) = \hat{x}^*(x_1, N)$. The domain corresponding to a cross-section at some x_1 will be denoted $\Omega'(x_1)$. Since $\partial\Omega$ is a hyperplane in the neighborhood of x^*, hence first coordinate of $x^*(N)$ is fixed for all $N > N_0$ and equal to x_1^*. Therefore the integral $I'(N)$ can be represented

$$I'(N) = \int_{x_1^*}^{\partial\Omega'} I'(x_1, N)dx_1,$$

where $\partial\Omega'$ is other then x_1^* boundary point of Ω' on the axis of coordinate x_1 and

$$I'(x_1, N) := \int_{\Omega(x_1)} g(x_1, \hat{x})e^{Nf(x_1,\hat{x},N)}d\hat{x}.$$

Since within Ω' property (8) is valid we can apply Theorem 2 for the above integral

$$I'(x_1, N) = e^{Nf(x_1,\hat{x}^*(x_1,N),N)}\left(\frac{2\pi}{N}\right)^{\frac{m}{2}}\left(\frac{g(x_1, \hat{x}^*(x_1, N))}{\sqrt{\det D^2 f(x_1, \hat{x}^*(x_1, N), N)}} + \frac{\omega_I(x_1, N)}{\sqrt{N}}\right).$$

Since we integrated over the set $\Omega(x_1)$ the constants which are in the bound of $\omega_I(x_1, N)$ can be bounded by the constants for larger set Ω listed in (2-7). Therefore $\omega_I(x_1, N)$ has the same bound as in the Theorem 2.
Now we integrate above expression w.r.t. x_1 over the set $(x_1^*, \partial\Omega')$ to obtain $I'(N)$. Since g divided by square root of determinant of considered Hessian matrix is bounded and has bounded derivative in the domain of integration we can use Theorem 1 which yields

$$I'(N) = e^{Nf(x^*(N),N)}\frac{1}{N}\left(\frac{2\pi}{N}\right)^{\frac{m}{2}}\left(\frac{g(x^*(N))}{|f'(x^*(N), N)|\sqrt{\det D^2 f(x^*(N), N)}} + \frac{\omega_{B1}(N)}{N} + \frac{\omega_I'(N)}{\sqrt{N}}\right),$$

where $(x_1^*, \hat{x}^*(x_1^*, N)) = x^*(N)$ and

$$|\omega_I'(N)| \leq |\omega_I(N)|\left(\frac{1}{|f'(x^*(N), N)|} + \frac{|\omega_{B2}(N)|}{N}\right),$$

Since Theorem 1 was applied on the domain which was the curve $\hat{x}^*(N)$ the constants which are in the bound of $\omega_{B1}(N)$ and $\omega_{B2}(N)$ can be bounded by the constants for larger set Ω listed in (2-7). Since the function g in the application of the Theorem 1 was equal to $g(x_1, \hat{x}^*(x_1, N))/\sqrt{\det D^2 f(x_1, \hat{x}^*(x_1, N), N)}$ and 1, respectively for ω_{B1} and ω_{B2} therefore we have estimates

$$\begin{aligned}|\omega_{B1}(N)| \leq & \frac{|g(x^*(N))|F^{(2)}}{\sqrt{\lambda}\left(F'^{(1)}\right)^3}e^{\frac{1}{2}F^{(2)}} + \frac{1}{\left(F'^{(1)}\right)^2}\left(\frac{G^{(1)}}{\sqrt{\lambda}} + \frac{mGF^{(3)}}{\lambda^{3/2}F'^{(2)}}\right) + \frac{G}{\sqrt{\lambda}F_{\Omega}'^{(1)}}Ne^{-N^{1/2}F_{\Omega}'^{(1)}} + \\ & + \frac{|g(x^*(N))|}{\sqrt{\lambda}F'^{(1)}}Ne^{-N^{1/2}F'^{(1)}}, \\ |\omega_{B2}(N)| \leq & \frac{F^{(2)}}{\left(F'^{(1)}\right)^3}e^{\frac{1}{2}F^{(2)}} + \frac{1}{F_{\Omega}'^{(1)}}Ne^{-N^{1/2}F_{\Omega}'^{(1)}} + \frac{1}{F'^{(1)}}Ne^{-N^{1/2}F'^{(1)}}.\end{aligned}$$

Then we combine above results with the estimate of $I''(N)$ and obtain statement of the theorem. $\square$

3 Probability application

We consider an integral (1) with additionally

$$f(x,N) = f(x) + \epsilon(N)\sigma(x), \tag{11}$$

where $\sigma(x)$ is some twice differentiable function at x^* and $\epsilon(N) \to 0$ as $N \to \infty$.
Then we have following result

Lemma 1. *The points of maximums of the functions in (11) are related by*

$$x^*(N) = x^* + \epsilon(N)O(1), \text{ as } N \to \infty, \tag{12}$$

Proof. The proof for the maximum of type (a) starts with use of the first order Taylor expansion

$$Df(x^*(N),N) = Df(x^*,N) + D^2f(x_\theta(N),N)^T(x^*(N) - x^*)$$

Since $x^*(N), x^*$ are a maximums of $f(x,N), f(x)$ respectively and by (11)

$$0 = \epsilon(N)D\sigma(x^*) + D^2f(x_\theta(N),N)^T(x^*(N) - x^*)$$

and then using fact that $|x - x^*|\|D^2f(x_\theta)^{-1}\|^{-1} \leq \left|D^2f(x_\theta)^T(x - x^*)\right|$ and (3) yields

$$\epsilon(N)\frac{|D\sigma(x^*)|}{F'^{(2)}} \geq |x^*(N) - x^*|,$$

which is the desired result.
For the maximum of type (b), as $\partial\Omega$ near x^* is a hyperplane, with appropriate rotation of coordinates if necessary, the first coordinate axis is orthogonal to that hyperplane. Therefore x_1 will be fixed for the points of maximum $x^*(N), N > N_0$ and in the other coordinates x^* and $x^*(N)$ are critical points. Therefore, the procedure of estimation is the same as for the previous type of maximum and yields the same result. $\square$

For every $N > N_0$ we define a probability measure on a measurable space $(\Omega, \mathcal{B}(\Omega))$

$$P_N(A) = \frac{\int_A e^{Nf(x,N)}dx}{\int_\Omega e^{Nf(x,N)}dx},$$

and corresponding random vector $X(N)$ defined as an identity map on Ω.

3.1 Weak Law of large numbers and Central Limit Theorem

Theorem 4 (Weak Law of large numbers)**.** *As $N \to \infty$ the random vector $X(N)$ converges in distribution to a constant vector x^* and we have following estimate of its mgf*

$$M_{X(N)}(\xi) = e^{\xi^T x^*}\left(1 + \frac{O(1)}{\sqrt{N}} + O(1)\epsilon(N)\right), \text{ as } N \to \infty.$$

Proof. We prove the theorem by showing convergence of the mgf of $X(N)$

$$M_{X(N)}(\xi) = \frac{\int_\Omega e^{\xi^T x} e^{Nf(x,N)} dx}{\int_\Omega e^{Nf(x,N)} dx}.$$

for $|\xi| < h$, for some $h > 0$.
With Theorem 2 we approximate the denominator for the two cases of maximum

$$\begin{aligned}
(a) \int_\Omega e^{Nf(x,N)} dx =& e^{Nf(x^*(N),N)} \left(\frac{2\pi}{N}\right)^{\frac{m}{2}} \frac{1}{\sqrt{\det D^2 f(x^*(N),N)}} \left(1 + \frac{O(1)}{\sqrt{N}}\right), \\
(b) \int_\Omega e^{Nf(x,N)} dx =& e^{Nf(x^*(N),N)} \frac{1}{N} \left(\frac{2\pi}{N}\right)^{\frac{m-1}{2}} \frac{1}{|f'(x^*(N),N)| \sqrt{|\det D^2 f(x^*(N),N)|}} \left(1 + \frac{O(1)}{\sqrt{N}}\right),
\end{aligned}$$

as $N \to \infty$.
Then we approximate the numerator, where the additional function in the exponent is g in the Theorems

$$\begin{aligned}
(a) \int_\Omega e^{\xi^T x + Nf(x,N)} dx =& e^{\xi^T x^*(N) + Nf(x^*(N),N)} \left(\frac{2\pi}{N}\right)^{\frac{m}{2}} \frac{1}{\sqrt{\det D^2 f(x^*(N),N)}} \left(1 + \frac{O(1)}{\sqrt{N}}\right), \\
(b) \int_\Omega e^{\xi^T x + Nf(x,N)} dx =& e^{\xi^T x^*(N) + Nf(x^*(N),N)} \frac{1}{N} \left(\frac{2\pi}{N}\right)^{\frac{m-1}{2}} \frac{1}{|f'(x^*(N),N)| \sqrt{|\det D^2 f(x^*(N),N)|}} \times \\
& \times \left(1 + \frac{O(1)}{\sqrt{N}}\right), \text{ as } N \to \infty.
\end{aligned}$$

Combining the approximations of denominators and numerators yields

$$M_{X(N)}(\xi) = e^{\xi^T x^*(N)} \left(1 + \frac{O(1)}{\sqrt{N}}\right), \text{ as } N \to \infty.$$

Next, we use estimate (12) and Taylor expansion for the exponent function to obtain

$$M_{X(N)}(\xi) = e^{\xi^T x^*} \Big(1 + \epsilon(N) \xi^T O(1)\Big) \left(1 + \frac{O(1)}{\sqrt{N}}\right),$$

which yields the result of the theorem. ☐

We consider a random vector $Y(N)$ for the two types of maximum

$$\begin{aligned}
&(a)\ Y(N) = \sqrt{N}(x^* - X(N)), \\
&(b)\ Y(N) = (N(x_1^* - X_1(N)), \sqrt{N}(\hat{x}^* - \hat{X}(N)))^T,
\end{aligned}$$

where $X(N) = (X_1(N), \hat{X}(N))$ and $x^* = (x_1^*, \hat{x}^*)$. For the case (b), in order to have $\partial\Omega$ at x^* orthogonal to axis of x_1 we might have to rotate the coordinate system with $Y(N)$ defined as inversely rotated.

Theorem 5 (Fluctuations theorem)**.** *As $N \to \infty$, if in (11) the function $\epsilon(N) = o\big(\frac{1}{\sqrt{N}}\big)$, then the distribution of $Y(N)$ converges weakly to a $\mathcal{N}(0, D^2 f(x^*)^{-1})$ distribution for the* *maximum of type (a). For the type (b) it converges to distribution with marginal distributions* $\exp\left(|f'(x^*)|\right)$ *and* $\mathcal{N}(0, D^2 f(x^*)^{-1})$ *for* $Y_1(N)$ *and* $(Y_2(N), \dots, Y_m(N))$ *respectively. Furthermore, as* $N \to \infty$ *we have following approximations of the mgf*

$$(a)\ M_{Y(N)}(\xi) = e^{\frac{1}{2}\xi^T D^2 f(x^*)^{-1}\xi}\left(1 + \frac{O(1)}{\sqrt{N}} + \epsilon(N)\sqrt{N}O(1)\right),$$

$$(b)\ M_{Y(N)}(\xi) = \frac{|f'(x^*)|}{|f'(x^*) - \xi_1|} e^{\frac{1}{2}\hat{\xi}^T D^2 f(x^*)^{-1}\hat{\xi}}\left(1 + \frac{O(1)}{\sqrt{N}} + \epsilon(N)\sqrt{N}O(1)\right),$$

where in the case (b) $\hat{\xi} = (\xi_2, \dots, \xi_m)$*,* $f'(x^*)$ *is a derivative w.r.t.* x_1 *and* $D^2 f(x^*)^{-1}$ *is a matrix of derivatives w.r.t. all but first coordinate.*

Proof. We define $\widetilde{f}(x, N) := f(x, N) + \frac{1}{N}\xi^T Y(N)$ and the mgf of $Y(N)$ is

$$M_{Y(N)}(\xi) = \frac{\int_\Omega e^{N\tilde{f}(x,N)}dx}{\int_\Omega e^{Nf(x,N)}dx}.$$

From the structure of $Y(N)$ we infer that for large enough N the properties of the function $\widetilde{f}$ are the same as of f but with the maximum at $\widetilde{x}^*(N)$.
First, we prove the result for the maximum of type (a). We approximate the numerator and denominator of the above mgf using Theorem 2 and obtain

$$M_{Y(N)}(\xi) = e^{N\widetilde{f}(\widetilde{x}^*(N),N) - Nf(x^*(N),N)} \frac{\sqrt{\det D^2 f(x^*(N), N)}}{\sqrt{\det D^2 \widetilde{f}(\widetilde{x}^*(N), N)}}\left(1 + \frac{O(1)}{\sqrt{N}}\right), \text{ as } N \to \infty.$$

Next, we insert the estimates (14) and (15) from Preposition 1

$$M_{Y(N)}(\xi) = e^{\frac{1}{2}\xi^T D^2 f(x^*)^{-1}\xi + N^{-1/2}O(1) + \epsilon(N)\sqrt{N}O(1)}\left(1 + \frac{O(1)}{\sqrt{N}}\right)\left(1 + \frac{O(1)}{\sqrt{N}}\right),$$

We estimate the second term in the exponent $\exp(N^{-1/2}O(1) + \epsilon(N)\sqrt{N}O(1)) = 1 + N^{-1/2}O(1) + \epsilon(N)\sqrt{N}O(1)$ and simplify whole estimate by including only expressions least decreasing in N and we get the first result of the theorem.
For the maximum of type (b) we approximate the numerator and denominator of the mgf using Theorem 3

$$M_{Y(N)}(\xi) = e^{N\widetilde{f}(\widetilde{x}^*(N),N) - Nf(x^*(N),N)} \frac{|f'(x^*(N), N)|}{|\widetilde{f}'(\widetilde{x}^*(N), N)|} \frac{\sqrt{\det D^2 f(x^*(N), N)}}{\sqrt{\det D^2 \widetilde{f}(\widetilde{x}^*(N), N)}}\left(1 + \frac{O(1)}{\sqrt{N}}\right),$$

where $\hat{x} = (x_2, \dots, x_m)$.
Since the boundary $\partial\Omega$ is a hyperplane near x^* and is orthogonal to axis of x_1 we have $\widetilde{x}_1^*(N) = x_1^*(N) = x_1^*$ and w.r.t. other coordinates we have maximum of type (a). Therefore for the above estimate we can apply results of Preposition 1 and obtain

$$M_{Y(N)}(\xi) = \frac{|f'(x^*(N), N)|}{|f'(\widetilde{x}^*(N), N) - \xi_1|} e^{\frac{1}{2}\hat{\xi}^T D^2 f(x^*)^{-1}\hat{\xi}}\left(1 + \frac{O(1)}{\sqrt{N}} + \epsilon(N)\sqrt{N}O(1)\right),$$

Next, we use first order Taylor theorem for $f'(\widetilde{x}^*(N), N)$ to obtain

$$|f'(\widetilde{x}^*(N), N) - f'(x^*, N)| \le F^{(2)}\big(|\widetilde{x}^*(N) - x^*(N)| + |x^*(N) - x^*|\big).$$

With use of (12), (13) and (11) we conclude

$$f'(\widetilde{x}^*(N),N) = f'(x^*) + \frac{O(1)}{\sqrt{N}}, \text{ as } N \to \infty.$$

Analogical estimation procedure is for $f'(x^*(N),N)$, with the result

$$f'(\widetilde{x}^*(N),N) = f'(x^*) + O(1)\epsilon(N), \text{ as } N \to \infty.$$

With above estimates we can express

$$\frac{|f'(x^*(N),N)|}{|f'(\widetilde{x}^*(N),N) - \xi_1|} = \frac{|f'(x^*)|}{|f'(x^*) + O(1)/\sqrt{N} - \xi_1|}|1 + \epsilon(N)O(1)| =$$
$$= \frac{|f'(x^*)|}{|f'(x^*) - \xi_1|}(1 + \epsilon(N)O(1))\left(\frac{1}{1 + \frac{O(1)}{|f'(x^*)-\xi_1|\sqrt{N}}}\right),$$

as $N \to \infty$ and substituting that into last estimate of mgf yields the final result. ☐

3.2 Additional estimates

Preposition 1. *For the defined previously functions f and $\widetilde{f}$ with the maximum of type (a) and provided that $\epsilon(N) = o(1/\sqrt{N})$ as $N \to \infty$ we have following approximations*

$$\widetilde{x}^*(N) - x^*(N) = D^2 f(x^*)^{-1}\frac{\xi}{\sqrt{N}} + \frac{O(1)}{N}, \tag{13}$$

$$\widetilde{f}(\widetilde{x}^*(N),N) - f(x^*(N),N) = \frac{1}{2N}\xi^T D^2 f(x^*)^{-1}\xi + \frac{O(1)}{N^{3/2}} + \frac{O(1)\epsilon(N)}{\sqrt{N}}, \tag{14}$$

$$\frac{\sqrt{\det D^2 f(x^*(N),N)}}{\sqrt{\det D^2 \widetilde{f}(\widetilde{x}^*(N),N)}} = 1 + \frac{O(1)}{\sqrt{N}}, \tag{15}$$

as $N \to \infty$.

Proof. We begin with the proof of the first estimate.
We use first order Taylor's expansion for the vector function $D\widetilde{f}(\widetilde{x}^*(N),N)$ at $x^*(N)$ and since $\widetilde{x}^*(N)$ and $x^*(N)$ are critical points

$$D^2 f(x_{\theta_1}(N),N)(\widetilde{x}^*(N) - x^*(N)) = \frac{1}{\sqrt{N}}\xi.$$

Since we have that $\widetilde{x}^*(N) \to x^*$ as $N \to \infty$, hence for large enough N the point $x_{\theta_1}(N)$ is in Ω' and since (3) consequently

$$\left|\widetilde{x}^*(N) - x^*(N)\right| \le \frac{(F'^{(2)})^{-1}|\xi|}{\sqrt{N}}$$

Further, we use second order Taylor's expansion of $D\widetilde{f}(\widetilde{x}^*(N),N)$ at $x^*(N)$ and with (17) we get an upper bound

$$\left|D^2 f(x^*(N),N)(\widetilde{x}^*(N) - x^*(N)) - \frac{1}{\sqrt{N}}\xi\right| \le \frac{F^{(3)}}{2}\left|\widetilde{x}^*(N) - x^*(N)\right|^2,$$

and substitute previous estimate in the RHS to obtain

$$\left|(\widetilde{x}^*(N)-x^*(N))-D^2f(x^*(N),N)^{-1}\frac{1}{\sqrt{N}}\xi\right|\leq\frac{F^{(3)}(F'^{(2)})^{-3}|\xi|^2}{2N}.$$

Next we apply Taylor's theorem at x^* to $D^2f(x^*(N),N)^{-1}$ and conclude with an esimate

$$\left\|D^2f(x^*(N),N)^{-1}-D^2f(x^*,N)^{-1}\right\|\leq\epsilon(N)F^{(3)}|O(1)|.$$

Since (11) and $(I+\epsilon A)^{-1}=I+O(\epsilon),$ as $\epsilon\to 0$

$$D^2f(x^*,N)^{-1}=D^2f(x^*)^{-1}\Big[I+\epsilon(N)D^2f(x^*)^{-1}\Big]^{-1}=D^2f(x^*)^{-1}+\epsilon(N)O(1),$$

Putting together above estimates we conclude

$$\left|(\widetilde{x}^*(N)-x^*(N))-\frac{1}{\sqrt{N}}D^2f(x^*)^{-1}\xi\right|\leq\frac{F^{(3)}(F'^{(2)})^{-3}|\xi|^2}{2N}+\frac{\epsilon(N)|O(1)\xi|}{\sqrt{N}},$$

and since $\epsilon(N)=o\big(\frac{1}{\sqrt{N}}\big)$

$$\left|(\widetilde{x}^*(N)-x^*(N))-\frac{1}{\sqrt{N}}D^2f(x^*)^{-1}\xi\right|\leq\frac{K_1}{N},$$

for some positive K_1. Hence we get the first result of the Preposition.
We start the proof of the second result with evaluating $\widetilde{f}(\widetilde{x}^*(N),N)$ with use of the above estimate

$$\widetilde{f}(\widetilde{x}^*(N),N)=f\bigg(x^*(N)+D^2f(x^*)^{-1}\frac{\xi}{\sqrt{N}}+\frac{O(1)}{N},N\bigg)-$$
$$-\frac{1}{\sqrt{N}}\xi^T\bigg(x^*(N)+D^2f(x^*)^{-1}\frac{\xi}{\sqrt{N}}+\frac{O(1)}{N}-x^*\bigg),$$

and expand f using 3-rd order Taylor expansion at $x^*(N)$ and then substitute into above

$$\widetilde{f}(\widetilde{x}^*(N),N)=f(x^*(N),N)-\frac{1}{\sqrt{N}}\xi^T\bigg(x^*(N)+D^2f(x^*)^{-1}\frac{\xi}{\sqrt{N}}+\frac{O(1)}{N}-x^*\bigg)+$$
$$+\frac{1}{2}\bigg(D^2f(x^*)^{-1}\frac{\xi}{\sqrt{N}}+\frac{O(1)}{N}\bigg)D^2f(x^*(N),N)\bigg(D^2(x^*)^{-1}\frac{\xi}{\sqrt{N}}+\frac{O(1)}{N}\bigg)+$$
$$+\frac{1}{6}D^{\otimes 3}f(x_{\theta_2}(N),N)\bigg(D^2f(x^*)^{-1}\frac{\xi}{\sqrt{N}}+\frac{O(1)}{N}\bigg)^{\otimes 3},$$

and get and upper bound as in (17)

$$\bigg|\widetilde{f}(\widetilde{x}^*(N),N)-f(x^*(N),N)+\frac{1}{\sqrt{N}}\xi^T\bigg(D^2f(x^*)^{-1}\frac{\xi}{\sqrt{N}}+\frac{O(1)}{N}+\epsilon(N)O(1)\bigg)-$$
$$-\frac{1}{2}\bigg(D^2f(x^*)^{-1}\frac{\xi}{\sqrt{N}}+\frac{O(1)}{N}\bigg)D^2f(x^*(N),N)\bigg(D^2f(x^*)^{-1}\frac{\xi}{\sqrt{N}}+\frac{O(1)}{N}\bigg)\bigg|\leq$$
$$\leq\frac{1}{6}F^{(3)}\bigg|D^2f(x^*)^{-1}\frac{\xi}{\sqrt{N}}+\frac{O(1)}{N}\bigg|^3.$$

Then we use first order Taylor's Theorem on $D^2f(x^*(N),N)$ at x^* and conclude with an estimate

$$D^2f(x^*(N),N) = D^2f(x^*,N) + \epsilon(N)O(1) = D^2f(x^*) + \epsilon(N)O(1), \tag{16}$$

where the second equality is due to (11).
We combine above estimates, apply Triangle and Schwarz inequalities to obtain

$$\begin{aligned}&\left|\widetilde{f}(\widetilde{x}^*(N),N) - f(x^*(N),N) + \frac{1}{2N}\xi^T D^2f(x^*)^{-1}\xi\right| \leq \frac{|O(1)\xi^T\epsilon(N)|}{\sqrt{N}} + \\ &+ \frac{|O(1)|}{N^{3/2}}\left[\frac{1}{6}F^{(3)}\left|D^2f(x^*)^{-1}\xi + \frac{O(1)}{\sqrt{N}}\right|^3 + \frac{1}{2}O(1)\epsilon(N)\sqrt{N}\left|D^2f(x^*)^{-1}\xi + \frac{O(1)}{\sqrt{N}}\right|^2 + \right. \\ &\left. + |\xi^T 1| + \frac{1}{2\sqrt{N}}\left\|D^2f(x^*)\right\|\right].\end{aligned}$$

Hence we have

$$\left|\widetilde{f}(\widetilde{x}^*(N),N) - f(x^*(N),N) + \frac{1}{2N}\xi^T D^2f(x^*)^{-1}\xi\right| \leq \frac{K_{21}}{N^{3/2}} + \frac{K_{22}\epsilon(N)}{\sqrt{N}},$$

where K_{21}, K_{22} are some positive constants, hence we get the second result.
Now we prove the third result of the Preposition. We start by taking first order Taylor approximation of $D^2\widetilde{f}(\widetilde{x}^*(N),N)$ at x^* and with use of (12) and the first result of this Preposition we obtain

$$\left\|D^2\widetilde{f}(\widetilde{x}^*(N),N) - D^2f(x^*,N)\right\| \leq \frac{F^{(3)}}{\sqrt{N}}\left[D^2f(x^*)^{-1}\xi + \frac{|O(1)|}{\sqrt{N}}\right].$$

Then, using property (11) yields

$$D^2\widetilde{f}(\widetilde{x}^*(N),N) = D^2f(x^*) + \frac{O(1)}{\sqrt{N}}.$$

Next, we divide estimate (16) by the above and get

$$\frac{D^2f(x^*(N),N)}{D^2\widetilde{f}(\widetilde{x}^*(N),N)} = I + \frac{O(1)}{\sqrt{N}}.$$

Further, the Taylor exapnsion of $f(x) = \sqrt{x}$ at 1 is

$$\sqrt{x} = 1 + (c)^{-1/2}(x-1),$$

where $c \in (\min(x,1), \max(x,1))$. We apply this expansion with

$$x = \det\left(\frac{D^2f(x^*(N),N)}{D^2\widetilde{f}(\widetilde{x}^*(N),N)}\right),$$

to obtain

$$\left|\frac{\sqrt{\det D^2f(x^*(N),N)}}{\sqrt{\det D^2\widetilde{f}(\widetilde{x}^*(N),N)}} - 1\right| = \frac{1}{\sqrt{c}}\left|\det\left(\frac{D^2f(x^*(N),N)}{D^2\widetilde{f}(\widetilde{x}^*(N),N)}\right) - 1\right|.$$

By (8) the determinants are bounded, therefore c is also bounded. Then with use Jacobi formula

$$\left|\det\left(\frac{D^2 f(x^*(N),N)}{D^2\widetilde{f}(\widetilde{x}^*(N),N)}\right)-1\right| = \left|\det\left(I+\frac{O(1)}{\sqrt{N}}\right)-1\right| \le \frac{K_3}{\sqrt{N}}.$$

where $K_3 > 0$. Hence, combining above results leads to a statement of the Preposition.

$\square$

A Appendix

For the the third term in the Taylor's theorem of a real function we have

$$\left|\sum_{i_1,i_2,i_3=1}^{3}\frac{f^3(x_\theta)}{\partial x_{i_1}\partial x_{i_2}\partial x_{i_3}}(x_{i_1}-x^*_{i_1})(x_{i_2}-x^*_{i_2})(x_{i_3}-x^*_{i_3})\right| = |D^{\otimes 3}f(x_\theta),(x-x^*)^{\otimes m}| =$$
$$= |\langle D^{\otimes 3}f(x_\theta),(x-x^*)^{\otimes m}\rangle| \le \|D^{\otimes 3}f(x_\theta)\|\|(x-x^*)^{\otimes m}\| = \|D^{\otimes 3}f(x_\theta)\||x-x^*|^m \quad (17)$$

where the last equality is due to the definition of tensor product.

Lemma 2. *For the integrer $N>0$, some constant $a>0$ we have*

$$\int_{\mathbb{R}^m\setminus S(N^{-1/3})}|x|^k e^{-aNx^2}dx \le \frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})}(aN)^{-\frac{k+m+1}{2}}\left(\Gamma(k+m)+\left(1+\sqrt{a}N^{1/6}\right)^{k+m-1}\right)e^{-aN^{1/3}}, \tag{18}$$

$$\int_{\mathbb{R}^m\setminus S(R)}|x|^k e^{-aNx^2}dx \le \frac{\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})}(aN)^{-\frac{k+m+1}{2}}\left(\Gamma(k+m)+\left(1+\sqrt{aN}R\right)^{k+m-1}\right)e^{-aNR^2}, \tag{19}$$

where k is a nonnegative integer and $S(N^{-1/3}), S(R)$ are m-dimensional spheres with the center at the origin and radius $N^{-1/3}$ and R respectively.

Proof. For the first integral we change the coordinate system into spherical, with $x = rs_r$, and radius $r = |x|$

$$\int_{\mathbb{R}^m\setminus S(N^{-1/3})}|x|^k e^{-aN|x|^2}dx = \int_{N^{-1/3}}^{\infty}\int_{S(r)} r^k e^{-aNr^2}ds_r dr.$$

As the function under the integral does not depend on the surface coordinates we can independently integrate over it. The surface of m-dimensional sphere of radius r is given by

$$S(r) = \frac{2\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})}r^{m-1},$$

and

$$\int_{N^{-1/3}}^{\infty}\int_{S(r)} r^k e^{-aNr^2}ds_r dr = \frac{2\pi^{\frac{m}{2}}}{\Gamma(\frac{m}{2})}\int_{N^{-1/3}}^{\infty} r^{k+m-1}e^{-aNr^2}dr.$$

Next, we perform a substitution $t = aN(r^2 - N^{-2/3})$ which yields

$$\int_{N^{-1/3}}^{\infty} r^{k+m-1}e^{-aNr^2}dr = \frac{1}{2}e^{-aN^{1/3}}(aN)^{-\frac{k+m+1}{2}}\int_0^{\infty}e^{-t}\left(t+aN^{1/3}\right)^{\frac{k+m-1}{2}}dt,$$

and we estimate the function $u(t) = (t + aN^{1/3})^{\frac{k+m-1}{2}}$ with $w(t) = t^{k+m-1}$ plus some constant. Let us calculate the points t when $u(t) = w(t)$. We solve this by reduction to quadratic equation with two roots

$$t_1 = \frac{1+\sqrt{1+4aN^{1/3}}}{2},\ t_2 = \frac{1-\sqrt{1+4aN^{1/3}}}{2},$$

Since $t_2 < 0$, quadratic function is convex and both functions are strictly increasing for $t \in [0, t_1)$ we have $u(t) > w(t)$ and opposite for $t \geq t_1$. Hence, the function $w(t) + w(t_1) \geq u(t)$ for all $t \geq 0$. Therefore we have an estimate

$$(t + aN^{\frac{1}{3}})^{(k+m-1)/2} \leq t^{k+m-1} + \left(1 + \sqrt{a}N^{1/6}\right)^{k+m-1}.$$

We substitute above into previous one and get that the integral is bounded

$$\int_0^\infty e^{-t}\left(t + aN^{1/3}\right)^{\frac{k+m-1}{2}} dt \leq \Gamma(k+m) + \left(1 + \sqrt{a}N^{1/6}\right)^{k+m-1},$$

hence putting the above results together gives the first result of Lemma.
The second result can be proved analogically. ☐

References

K.W. Breitung. *Asymptotic Approximations for Probability Integral.* Springer, 1994.

Vassili N. Kolokoltsov. *Semiclassical Analysis for Diffusions and Stochastic processes.* Springer, 2000.

R. Wong. *Asymptotic Approximations for Integrals.* SIAM, 2001.